

Department of Computer Science,
University of Otago

UNIVERSITY
of
OTAGO



Te Whare Wānanga o Ōtāgo

Technical Report OUCS-2002-10

Restricted permutations and queue jumping

Authors:

**M.H. Albert, M.D. Atkinson, H. van
Ditmarsch, C.C. Handley**
Department of Computer Science

R.E.L. Aldred, D.A. Holton
Department of Mathematics and Statistics

Status: Submitted for publication



Department of Computer Science,
University of Otago, PO Box 56, Dunedin, Otago, New Zealand

<http://www.cs.otago.ac.nz/trseries/>

Restricted permutations and queue jumping

M. H. Albert* R. E. L. Aldred† M. D. Atkinson*
H. P. van Ditmarsch* C. C. Handley* D. A. Holton†

November 12, 2002

Abstract

A connection between permutations that avoid 4231 and a certain queueing discipline is established. It is proved that a more restrictive queueing discipline corresponds to avoiding both 4231 and 42513, and enumeration results for such permutations are given.

1 Introduction

Let $s_n(\alpha_1, \alpha_2, \dots)$ be the number of permutations of length n that avoid all of the permutation patterns $\alpha_1, \alpha_2, \dots$. Finding a formula or a generating function for $s_n(\alpha_1, \alpha_2, \dots)$ is a difficult and much studied problem. For a single pattern α the sequence $s_n(\alpha)$ is known only in the following cases (see [2, 3, 4]):

- $\alpha = 12 \dots k$ or $\alpha = k \dots 21$ for any k
- $|\alpha| \leq 3$
- $|\alpha| = 4$ but $\alpha \neq 1324, 4231$

So the first unsolved cases are $\alpha = 1324$ and $\alpha = 4231$ which are equivalent by symmetry. Here the lower bound $s_n(4231) \geq s_n(4321)$ has been proved by Bóna [1] who has also made some contributions towards an upper bound.

In this note we introduce a certain type of queue and, harking back to early work by Knuth [4] on such problems, study its connection with 4231-avoiding permutations. By strengthening the conditions satisfied by the queue we go on to solve the enumeration problem for the sequence $s_n(4231, 42513)$.

*Department of Computer Science, University of Otago

†Department of Mathematics and Statistics, University of Otago

2 Jump queues

We shall define two queue-based data structures. They will be used with an input sequence $1, 2, \dots, n$ whose members are added one by one to the rear of the queue. Removals from the queue generate an output sequence that will be a permutation of $1, 2, \dots, n$. Removals are always allowed from the front of the queue but both our queue-based structures also allow elements other than the front member of the queue to be output, that is *queue jumping* (when we refer to a *jump* we will always mean an output operation which would not be permitted by an ordinary queue). When a queue jump occurs some of the items in the queue become “locked” (forbidden to jump until the lock is released).

Our two structures differ in the extent to which instances of queue jumping restrict further queue jumping. The first of these data structures, the *loosely locked jump queue* is defined by the property that when an element x is jumped from the queue all the elements in the queue behind x become locked; they are released (given the freedom to jump) when all the elements in front of x have been output. Note that any new elements added to a loosely locked jump queue are initially not locked.

In the second structure, the *strictly locked jump queue*, the locking rule is more severe. Again, elements in the queue behind an element x that jumps become locked; furthermore, any new elements that are added to a queue that already has some locked elements, are initially in the locked state. As before, the lock on an element is released once all the elements in front of the jumped item that initially caused the lock have been output.

In both cases jumping from the rear of the queue imposes no locks so our queues are at least as powerful as the input-restricted dequeues that were analysed in [4].

In studying the output permutations generated by either type of queue we observe that an output permutation may be producible in several ways. This allows us to make a simplifying assumption. Suppose that we are attempting to generate a particular permutation π and have proceeded to a point where we wish to output a symbol p . If p is already in the queue then if it is possible to succeed at all from this point onwards, we can succeed by outputting p immediately. For the only alternative is to add further elements to the queue and then output p . The only effect that this might have (versus outputting p immediately and then adding the same elements) is to lock some queue elements that would not be locked in the original instance. So, it cannot be harmful to do any output as soon as it becomes available and, from now on, we consider only operation sequences with this property. Under that assumption we will regard the production of any permutation π as taking place in a number of *stages*. In any of these stages one or more input elements are added to the queue, the last of these is then output (such outputs produce the left to right maxima of π), and then further output from the queue occurs (possibly none at all); a stage comes to an end when the next element of π to be output has not yet been added to the queue.

We begin our investigation with a result whose easy proof is omitted.

Lemma 1 *Suppose we have a jump queue of either sort with a frontal segment $\alpha = a_1 < a_2 < \dots < a_m$. Then the permutations of α that can be generated by queue removals (from the front or by jumping) are precisely those that avoid 231.*

Proposition 2 *The collection of permutations that can be produced by a loosely locked jump queue is the class of 4231-avoiding permutations.*

Proof: Let a permutation π be given which contains a 4231 pattern as

$$\pi = \dots d \dots b \dots c \dots a \dots$$

and suppose that we could produce π using a loosely locked jump queue. In order to output d before all of a , b , and c , those elements must be in the queue when d is output. However, the subsequent output of b would then lock c so that it could not be output until a was. So, in fact, π could not be generated.

Conversely, suppose that a permutation π avoids 4231. Let $m_1 < m_2 < \dots < m_k$ be the left to right maxima of π . Then we can write

$$\pi = m_1 \alpha_1 m_2 \alpha_2 \dots m_k \alpha_k$$

where each α_i is some segment of π and $m_i > \alpha_i$. For convenience define $m_0 = 0$.

We will show that π can be produced by following the operation of the queue in attempting to produce it, and observing that we never reach a point where an element which we need to output is locked. We argue inductively on the stages of this production (as defined previously) where stage j produces the segment $m_j \alpha_j$.

In the first stage the elements from 1 through m_1 are added to the queue, then m_1 is output (without causing any locks), and then the elements of α_1 must now be output. At present this can certainly be accomplished since α_1 avoids 231. However, this sequence of operations may leave some remaining elements of $[1, m_1)$ locked in the queue.

Suppose that, after j stages, we have output the initial segment $m_1 \alpha_1 \dots m_j \alpha_j$ of π . Stage $j + 1$ begins by adding the elements of $(m_j, m_{j+1}]$ to the queue and then outputting the element m_{j+1} . Next we begin to output the elements of α_{j+1} . Consider the point at which an element c of this type is to be output. Choose $i \leq j + 1$ such that $m_{i-1} < c < m_i$. Any lock to the output of c would have been applied by an element $b < c$ jumping after c had been added to the queue (i.e. after the output of m_i). For this lock to have remained in force there must be an element $a < b$ still in the queue. But if all this were true the elements $m_i b c a$ would form a 4231 pattern in π . Stage $j + 1$ therefore succeeds in producing a further segment $m_{j+1} \alpha_{j+1}$ of output and the inductive proof is complete. ■

Proposition 3 *The collection of permutations that can be produced by a strictly locked jump queue is the class of $\{4231, 42513\}$ -avoiding permutations.*

Proof: The proof is similar in spirit to that above, so we will provide somewhat fewer details. In one direction, suppose that π contains the pattern 42513 as the subsequence $dbeac$ and yet can be produced by the queue. Then the element c would be locked by the output of b . Since it is required that a be output after e the element a must still be in the queue when e enters and so c would still be locked at this point; therefore e would be locked upon entering the queue and could not be output at the proper place. Obviously if π contains 4231 it cannot be output, for even a loosely locked queue would not suffice in that case.

Conversely, suppose that π avoids these two patterns. Write

$$\pi = m_1 \alpha_1 m_2 \alpha_2 \cdots m_k \alpha_k$$

as above, and follow the operation of the queue in stages again.

Suppose that the output of m_j is prevented because it has been locked by some preceding jump of an element b . Then b was preceded by some $m_i > b$, and some element $c > b$ still remains in the queue as does some element $a < b$ (or m_j would no longer be locked). Then one of the two sequences $m_i b m_j c a$ or $m_i b m_j a c$ occurs in π . The latter is a 42513 pattern and the former contains a 4231 pattern, a contradiction in either case.

Now suppose that the output of some x in α_j is prevented by a lock caused by some preceding jump of an element b . Then choose m_i , c , and a as above. Since the pattern 4231 does not occur in π the elements m_i , b , x , a , c occur in π in that order. If $x > m_i$ we have a pattern 42513 while if $x < m_i$ then $m_i b x a$ is a 4231 pattern. Again, a contradiction is achieved in either case. ■

3 The number of permutations avoiding 4231 and 42513

Consider the operation of a strictly locked jump queue as it produces some permutation that avoids 4231 and 42513. At a point where no elements are locked we might choose to add one or more input elements (after which our next output step must be to remove the last element of the queue), to jump an element from the rear of the queue (which imposes no locks), or to output an earlier element, say the j th. In the latter case we must, or rather might as well, output all elements (if any) which are earlier still in the queue before continuing the operation. In so doing, we can produce any 231-avoiding permutation of these $j - 1$ elements. As is well known, the number of such permutations is c_{j-1} , the $(j - 1)$ st Catalan number.

If we set q to be the number of elements in the queue, and i the number remaining in the input then this trichotomy is easily translated into a recurrence for the

number of permutations that can be produced from this configuration. However, the manipulation of the resultant quantities will be simplified if we make a distinction between two cases: where the next output is the last element of the queue, or where we do not place any restriction on the next output. Let $l(q, i)$ enumerate the former class and $n(q, i)$ the latter (both quantities conventionally 0 if either q or i is negative). Then we have:

$$\begin{aligned} l(q, i) &= l(q+1, i-1) + n(q-1, i) \\ n(q, i) &= l(q, i) + \sum_{j=1}^{q-1} c_{j-1} n(q-j, i). \end{aligned}$$

Consider the power series:

$$\begin{aligned} N(x, t) &= \sum_{q, i \geq 0} n(q, i) x^q t^i \\ L(x, t) &= \sum_{q, i \geq 0} l(q, i) x^q t^i \\ C(x) &= \sum_{i \geq 0} c_i x^i \\ f(t) &= L(0, t), \end{aligned}$$

the last of which, by Proposition 3, is the generating function for the class of permutations that avoid 4231 and 42513. Then the recurrences translate easily into the equations:

$$\begin{aligned} L(x, t) &= \frac{t(L(x, t) - f(t))}{x} + xN(x, t) + 1 \\ N(x, t) &= L(x, t) + xN(x, t)C(x) - xf(t)C(x). \end{aligned}$$

After some rearrangement this yields:

$$L(x, t) ((x^2 - xt)C(x) + x^2 - x + t) = (t - xtC(x) + x^3C(x)) f(t) - x + x^2C(x).$$

Now the stage is set for a simple application of the kernel method. Consider the circumstances under which the parenthetical expression in the left hand side above is zero:

$$\begin{aligned} (x^2 - xt)C(x) + x^2 - x + t &= 0 \\ xC^2(x) - C(x) + 1 &= 0. \end{aligned}$$

Solving formally for $C(x)$ and x in terms of t yields:

$$\begin{aligned} C^3(x)t - C(x) + 1 &= 0 \\ x &= C(x)t \end{aligned}$$

These conditions can then be substituted in the right hand side, which must also be zero, yielding eventually:

$$f(t) = \frac{1}{1 - \eta(t)t}$$

where

$$t\eta(t)^3 - \eta(t) + 1 = 0.$$

The latter equation is easily seen to be the equation satisfied by the generating function for ternary trees. Therefore the n th term of the power series expansion of $\eta(t)$ is the number of ternary trees on n nodes, namely

$$\frac{\binom{3n}{n-1}}{n}$$

From this it follows readily that, if f_n denotes the n th term of $f(t)$,

$$\liminf_{n \rightarrow \infty} \sqrt[n]{f_n} = 27/4$$

and more detailed asymptotics could easily be obtained. From the above equations we can also read off the recurrence

$$f_n = \sum_{i=0}^{n-1} f_{n-1-i} \binom{3i}{i-1} / i$$

and thereby compute the expansion of

$$f(t) = 1 + t + 2t^2 + 6t^3 + 23t^4 + 102t^5 + 495t^6 + 2549t^7 + 13682t^8 + 75714t^9 + \dots$$

to as many terms as necessary.

References

- [1] M. Bóna: Permutations avoiding certain patterns, The case of length 4 and generalizations, *Discrete Math.* 175 (1997) 55–67.
- [2] M. Bóna: Exact enumeration of 1342-avoiding permutations, A close link with labeled trees and planar maps, *J. Combin. Theory, Ser. A*, 80 (1997) 257–272.
- [3] I. M. Gessel: Symmetric functions and P -recursiveness, *J. Combin. Theory Ser. A*, 53 (1990), 257–285.
- [4] D.E. Knuth: *Fundamental Algorithms, The Art of Computer Programming* Vol. 1 (Second Edition), Addison-Wesley, Reading, Mass. (1973).