

Sorting classes

M. H. Albert

Department of Computer Science
University of Otago
`malbert@cs.otago.ac.nz`

R. E. L. Aldred

Department of Mathematics and Statistics
University of Otago
`raldred@math.otago.ac.nz`

M. D. Atkinson

Department of Computer Science
University of Otago
`mike@cs.otago.ac.nz`

C. C. Handley

Department of Computer Science
University of Otago
H. van Ditmarsch

Department of Computer Science
University of Otago
`hans@cs.otago.ac.nz`

D. A. Holton

Department of Mathematics and Statistics
University of Otago
`dholton@math.otago.ac.nz`

D. J. McCaughan

Department of Mathematics and Statistics
University of Otago
`dmccaughan@math.otago.ac.nz`

Submitted: Dec 20, 2004; Accepted: June 17, 2005; Published: June 30, 2005
Mathematics Subject Classifications: 05A15, 05A16

Abstract

Weak and strong sorting classes are pattern-closed classes that are also closed downwards under the weak and strong orders on permutations. They are studied using partial orders that capture both the subpermutation order and the weak or strong order. In both cases they can be characterised by forbidden permutations in the appropriate order. The connection with the corresponding forbidden permutations in pattern-closed classes is explored. Enumerative results are found in both cases.

1 Introduction

A permutation π is said to be a *subpermutation* of a permutation σ (or to be *involved* in σ) if σ has a subsequence that is ordered in the same relative way as π . For example 231 is a subpermutation of 35412 because of its subsequence 351 which has the same pattern as 231. We say that σ *avoids* π if π is not

a subpermutation of σ . The developing theory of permutation patterns is now a well-established part of combinatorics (see, for example, [12]).

This theory was originally motivated by the study of the sortable permutations associated with various computing devices (abstract data types such as stacks and deques [8], token passing networks [3], or hardware switches [2]). All these devices have the property that, if they are able to sort a sequence σ , then they are able to sort any subsequence of σ .

This *subsequence* property (that subsequences of sortable sequences are themselves sortable) is a very natural one to postulate of a sorting device. It is exactly this property that guarantees that the set of sortable permutations is closed under taking subpermutations. But there are other natural properties that a sorting device might have. We are particularly interested in the following two. Both of them reflect the idea that “more sorted” versions of sortable sequences should themselves be sortable.

1. If $s_1 s_2 \dots s_n$ is sortable and $s_i > s_{i+1}$ then $s_1 s_2 \dots s_{i-1} s_{i+1} s_i \dots s_n$ is sortable, and
2. If $s_1 s_2 \dots s_n$ is sortable and $s_i > s_j$ where $i < j$ then

$$s_1 s_2 \dots s_{i-1} s_j s_i s_{i+1} \dots s_{j-1} s_i s_{j+1} \dots s_n$$

is sortable.

For the moment we call these the *weak* and *strong exchange* properties (the second obviously implies the first). The weak exchange property would hold for sorting devices that operated by exchanging adjacent out of order pairs while the strong exchange property would hold if arbitrary out of order pairs could be exchanged. Our paper is about the interaction between each of these properties and the subsequence property.

We shall study this interaction using various (partial) orders on the set Ω of all (finite) permutations. Since we shall be considering several partial orders on Ω we shall write $\sigma \mathcal{P} \tau$ when we mean that $\sigma \leq \tau$ in the partial order \mathcal{P} ; this avoids the confusion of the symbol “ \leq ” being adorned by various subscripts. In the same spirit we write $\sigma \overline{\mathcal{P}} \tau$ to mean $\sigma \not\leq \tau$ in \mathcal{P} .

All the partial orders we study will satisfy the minimum condition (that is, all properly descending chains are finite) and we shall assume this from now on.

If \mathcal{P} is a partial order on Ω the *lower ideals* of \mathcal{P} are those subsets X of Ω with the property

$$\beta \in X \text{ and } \alpha \mathcal{P} \beta \implies \alpha \in X.$$

Since \mathcal{P} satisfies the minimum condition such a lower ideal can be studied through the set $b(X)$ of minimal permutations of $\Omega \setminus X$. Obviously $b(X)$ determines X uniquely since

$$X = \{\beta \mid \alpha \overline{\mathcal{P}} \beta \text{ for all } \alpha \in b(X)\}.$$

In the classical study of permutation patterns we use the subpermutation order that we denote by \mathcal{I} (standing for involvement). The lower ideals of \mathcal{I} are generally the central objects of study and are called *closed classes*. If X is a closed class then $b(X)$ is called the *basis* of X . Indeed the most common way of describing a closed class is by giving its basis (and therefore defining it by avoided patterns). We write $av(B)$ to denote the set of permutations which avoid all the permutations of the set B . If a closed class is not given in this way then, often, the first question is to determine the basis. A second question, perhaps of even greater interest, is to enumerate the class; in other words, to determine by formula, recurrence or generating function how many permutations it has of each length.

However, these questions can be posed for *any* partial order on Ω and much of our paper is devoted to answering them for orders that capture the subsequence property *and* the weak or strong exchange properties.

A closed class is called a *weak sorting class* if it has the weak exchange property and a *strong sorting class* if it has the strong exchange property.

Our aim is to set up a framework within which these two notions can be investigated and to exploit this framework by proving some initial results about them. We shall begin by investigating the two natural analogues of the subpermutation order that are appropriate for these two concepts. In particular there are natural notions of a basis for each type of sorting class; we shall explore how the basis of a sorting class is related to the ordinary basis and use this to derive enumerative results. In the remainder of this section we set up the machinery for studying sorting classes and then survey the main results of Sections 2 and 3 on weak and strong sorting classes respectively.

The terms ‘weak’ and ‘strong’ have been chosen to recall two important orders on the set of permutations of length n : the weak and strong orders. For completeness we shall give their definitions below (extended to the set Ω of permutations of all lengths). In these definitions and elsewhere in the paper we use Roman lower case letters for the individual symbols within a permutation and Greek lower case letters for sequences of zero or more symbols.

The *weak order* \mathcal{W} on Ω can be defined as the transitive closure of the set of pairs

$$\mathcal{W}_0 = \{(\lambda r s \mu, \lambda s r \mu) \mid r < s\}.$$

The *strong order* \mathcal{S} on Ω can be defined as the transitive closure of the set of pairs

$$\mathcal{S}_0 = \{(\lambda r \mu s \nu, \lambda s \mu r \nu) \mid r < s\}.$$

Notice that, for both \mathcal{W} and \mathcal{S} , only permutations of equal length can be comparable.

The subpermutation order \mathcal{I} on Ω can be defined as the transitive closure of the set of pairs

$$\mathcal{I}_0 = \{(\lambda \mu, \lambda' r \mu')\}$$

where $\lambda' \mu'$ is order isomorphic to $\lambda \mu$.

Weak (respectively, strong) sorting classes are the lower ideals in the partial order defined by the transitive closure of $\mathcal{I} \cup \mathcal{W}$ (respectively $\mathcal{I} \cup \mathcal{S}$) and so can be studied using the same machinery that has been used for arbitrary closed classes, adapted to the appropriate order.

We begin by giving a simple description of these transitive closures. In this description we denote the relational composition of two partial orders by juxtaposition.

Lemma 1 *The transitive closure of $\mathcal{I} \cup \mathcal{W}$ is $\mathcal{I}\mathcal{W}$ while that of $\mathcal{I} \cup \mathcal{S}$ is $\mathcal{I}\mathcal{S}$. In fact $\mathcal{W}\mathcal{I} = \mathcal{I}\mathcal{W}$ while $\mathcal{S}\mathcal{I}$ is strictly included in $\mathcal{I}\mathcal{S}$.*

Proof: Suppose that $\alpha \mathcal{I} \beta \mathcal{W}_0 \gamma$ represents a pair (α, γ) of the relation $\mathcal{I}\mathcal{W}_0$. Let $\alpha = a_1 a_2 \dots$ and let $a'_1 a'_2 \dots$ be a subsequence of β order isomorphic to α . Let xy be the two adjacent symbols of β that become yx in γ . If none or one of these is one of the a'_i then $\alpha \mathcal{I} \gamma$. If both of them are among $a'_1 a'_2 \dots$ then they must be a'_i and a'_{i+1} for some i . Let β' be the result of exchanging a_i and a_{i+1} in α ; then we have $\alpha \mathcal{W} \beta' \mathcal{I} \gamma$. This proves that $\mathcal{I}\mathcal{W}_0 \subseteq \mathcal{W}\mathcal{I}$ and it follows readily that $\mathcal{I}\mathcal{W}_0^t \subseteq \mathcal{W}\mathcal{I}$ for all t and hence that $\mathcal{I}\mathcal{W} \subseteq \mathcal{W}\mathcal{I}$.

To prove the opposite inclusion suppose that $\alpha \mathcal{W}_0 \beta \mathcal{I}_0 \gamma$ represents a pair (α, γ) of the relation $\mathcal{W}_0 \mathcal{I}_0$. Then we have

$$\begin{aligned} \alpha &= \theta a b \phi \\ \beta &= \theta b a \phi \end{aligned}$$

and γ is obtained from β by inserting an extra symbol x (with appropriate renumbering of the symbols larger than x).

If x does not occur between b and a then we can consider γ to be obtained from α by first inserting x and then swapping a and b ; so, in this case, $\alpha\mathcal{I}_0\mathcal{W}\gamma$. If x occurs between b and a then, depending on the value of x , we define ξ as either $\theta xab\phi, \theta axb\phi, \theta abx\phi$ so that the three symbols a, b, x come in increasing order. Then

$$\theta ab\phi \mathcal{I}_0 \xi \mathcal{W} \theta bxa\phi$$

and so, again, $\alpha \mathcal{I}_0 \mathcal{W} \gamma$.

We have proved that $\mathcal{W}_0\mathcal{I}_0 \subseteq \mathcal{I}_0\mathcal{W}$ and it readily follows that $\mathcal{W}\mathcal{I}_0 \subseteq \mathcal{I}_0\mathcal{W}$, and then that $\mathcal{W}\mathcal{I} \subseteq \mathcal{I}\mathcal{W}$. The transitive closure of $\mathcal{I} \cup \mathcal{W}$ is, by definition,

$$\bigcup_{i=0}^{\infty} (\mathcal{I} \cup \mathcal{W})^i.$$

However, \mathcal{I} and \mathcal{W} are transitively closed and $\mathcal{W}\mathcal{I} \subseteq \mathcal{I}\mathcal{W}$, and so this expression simplifies to $\mathcal{I}\mathcal{W}$.

Suppose now that $\alpha \mathcal{S}_0 \beta \mathcal{I} \gamma$ represents a pair (α, γ) of the relation $\mathcal{S}_0\mathcal{I}$. Put $\alpha = \lambda r \mu s \nu$ with $r < s$ and $\beta = \lambda s \mu r \nu$. Let $\lambda' s' \mu' r' \nu'$ denote a subsequence of γ order isomorphic to β . Consider the permutation γ' obtained from γ by interchanging s' and r' . Clearly $\alpha \mathcal{I} \gamma' \mathcal{S} \gamma$. This shows that $\mathcal{S}_0\mathcal{I} \subseteq \mathcal{I}\mathcal{S}$. But then it follows, as above, that $\mathcal{S}\mathcal{I} \subseteq \mathcal{I}\mathcal{S}$. However 321 \mathcal{I} 1432 \mathcal{S} 3412 yet there exists no permutation θ with 321 $\mathcal{S} \theta \mathcal{I}$ 3412; therefore the inclusion is strict.

It follows as above that $\mathcal{I}\mathcal{S}$ is the transitive closure of $\mathcal{I} \cup \mathcal{S}$. ■

The orders $\mathcal{I}\mathcal{W}$ and $\mathcal{I}\mathcal{S}$ have fewer symmetries (2 and 4 respectively) than the subpermutation order (which has 8). In the following elementary result, if $\zeta = z_1, \dots, z_n$, ζ^* denotes the ‘reverse complement’ of ζ

$$\zeta^* = n + 1 - z_n, n + 1 - z_{n-1}, \dots, n + 1 - z_1.$$

Lemma 2 *Let ξ, ζ be permutations. Then*

1. $\xi \mathcal{I}\mathcal{W} \zeta \iff \xi^* \mathcal{I}\mathcal{W} \zeta^*$, and
2. $\xi \mathcal{I}\mathcal{S} \zeta \iff \xi^{-1} \mathcal{I}\mathcal{S} \zeta^{-1} \iff \xi^* \mathcal{I}\mathcal{S} \zeta^*$.

We have already noted that every closed class X can be described by a forbidden pattern set T as

$$av(T) = \{\sigma \mid \beta \overline{\mathcal{I}} \sigma \text{ for all } \beta \in T\}.$$

We can describe weak and strong sorting classes in a similar way using the orders $\mathcal{I}\mathcal{W}$ and $\mathcal{I}\mathcal{S}$. In other words, given a set T of permutations we define

$$\begin{aligned} av(T, \mathcal{I}\mathcal{W}) &= \{\sigma \mid \beta \overline{\mathcal{I}\mathcal{W}} \sigma \text{ for all } \beta \in T\}. \\ av(T, \mathcal{I}\mathcal{S}) &= \{\sigma \mid \beta \overline{\mathcal{I}\mathcal{S}} \sigma \text{ for all } \beta \in T\}. \end{aligned}$$

which are weak and strong sorting classes respectively. Every weak and strong sorting class X can be defined in this way taking for T that set of permutations minimal with respect to $\mathcal{I}\mathcal{W}$ or $\mathcal{I}\mathcal{S}$ not belonging to X . If T is the minimal avoided set then it is tempting to call it the basis of the class it defines. Unfortunately that leads to a terminological ambiguity since both $av(T, \mathcal{I}\mathcal{W})$ and $av(T, \mathcal{I}\mathcal{S})$ are pattern closed classes and so have bases in the ordinary sense. To avoid such confusion we shall use the terms weak basis and strong basis. However, two significant questions now arise. If we have defined a weak sorting class by its weak basis, what is its basis in the ordinary sense? Similarly for strong sorting classes, what is the connection between the strong basis and the ordinary basis?

In the next section, on weak sorting classes, we shall see that the first of these questions has a relatively simple answer. In that section we also give a general result about the weak sorting class defined by a

basis that is the direct sum of two sets. We go on to enumerate weak sorting classes whose weak basis is a single permutation of length at most 4.

In the final section, on strong sorting classes, we shall see that the ordinary basis is not easily found from the strong basis. Nevertheless we can define a process that constructs the ordinary basis from the strong basis; and we prove that the ordinary basis is finite if the strong basis is finite. We have used this process as a first step in enumerating strong sorting classes defined by a single strong basis element of length at most 4. We shall give a summary of these results and some remarks on their proofs.

We also introduce a 2-parameter family of strong sorting classes denoted by $\mathcal{B}(r, s)$. These classes are important because every (proper) strong sorting class is contained in one (indeed infinitely many) of them. We shall show how the $\mathcal{B}(r, s)$ can be enumerated and give a structure theorem that expresses $\mathcal{B}(r, s)$ as a composition of very simple strong sorting classes.

2 Weak sorting classes

Proposition 3 *Let T be a set of permutations and let*

$$T' = \{\sigma \mid \tau \mathcal{W} \sigma \text{ for some } \tau \in T\}$$

(the upper weak closure of T). Then

$$av(T, \mathcal{I}\mathcal{W}) = av(T, \mathcal{W}\mathcal{I}) = av(T').$$

Proof: The first equality is immediate from Lemma 1. To prove the second, first suppose that $\sigma \notin av(T, \mathcal{W}\mathcal{I})$. Then, for some $\tau \in T$, we have $\tau \mathcal{W}\mathcal{I} \sigma$. Hence there exists $\tau' \in T'$ with $\tau \mathcal{W} \tau' \mathcal{I} \sigma$. The final relation says that $\sigma \notin av(T')$.

Conversely, suppose that $\sigma \notin av(T')$. Then, for some $\tau' \in T'$, we have $\tau' \mathcal{I} \sigma$. By definition of T' there exists $\tau \in T$ with $\tau \mathcal{W} \tau'$. But then $\tau \mathcal{W}\mathcal{I} \sigma$ which means that $\sigma \notin av(T, \mathcal{W}\mathcal{I})$. ■

Corollary 4 *The class $av(T)$ is a weak sorting class if and only if every permutation in the upward weak closure of T involves a permutation of T .*

Proof: Let T' be the upward weak closure of T . Then, by the previous proposition, $av(T, \mathcal{I}\mathcal{W}) = av(T')$ and so $av(T)$ is a weak sorting class if and only if $av(T) = av(T')$. The Corollary now follows. ■

Corollary 5 *If a weak sorting class has a finite weak basis then its ordinary basis is also finite.*

Proof: Let T be the weak basis of a weak sorting class and let T' be its upward weak closure. Obviously, T' is finite if T is finite. While T' may not be the ordinary basis of $av(T')$ (since it might not be an antichain) this ordinary basis just consists of the minimal elements of T' and so is finite. ■

To state the next result we need to recall the notion of the direct sum of two sets of permutations and some related terms. If α and β are permutations of lengths m and n then $\alpha \oplus \beta$ is the permutation of length $m + n$ whose first m symbols are all smaller than the last n symbols, the first m symbols comprise a sequence isomorphic to α , and the last n symbols comprise a sequence isomorphic to β . We extend this notion to sets X and Y of permutations by defining

$$X \oplus Y = \{\alpha \oplus \beta \mid \alpha \in X, \beta \in Y\}.$$

We also recall that a permutation is said to be *indecomposable* if it cannot be expressed as $\alpha \oplus \beta$. Every permutation has a unique expression in the form $\alpha_1 \oplus \cdots \oplus \alpha_k$ where each α_i is indecomposable, and the α_i are called the components of α . Closed classes whose basis elements are all indecomposable are somewhat easier to handle than arbitrary ones. This is because they have the property of being closed under direct sums and can be enumerated if their indecomposables can be enumerated [4].

Theorem 6 Let R, S be the weak bases of weak sorting classes \mathcal{A}, \mathcal{B} and let \mathcal{C} be the weak sorting class whose weak basis is $T = R \oplus S$. Let $(a_n), (b_n), (c_n)$ be the enumeration sequences for $\mathcal{A}, \mathcal{B}, \mathcal{C}$ and let $a(t), b(t), c(t)$ be the associated exponential generating functions. Then

$$c(t) = (t - 1)a(t)b(t) + a(t) + b(t).$$

Proof: Let R', S', T' be the upward weak closures of R, S, T . By Proposition 3, we have $\mathcal{A} = av(R')$, $\mathcal{B} = av(S')$, and $\mathcal{C} = av(T')$. We can compute the structure of the permutations of T' using the property that they are in the upward weak closure of some $\rho \oplus \sigma$ ($\rho \in R, \sigma \in S$). Such permutations must be the union of two sequences ρ', σ' where

1. $\rho' < \sigma'$, and
2. ρ', σ' are (order isomorphic to) permutations of R', S' .

Conversely, every such permutation is in the upward weak closure of some $\rho \oplus \sigma \in R \oplus S$ and so lies in T' .

From this description we can determine the structure of permutations in \mathcal{C} . We describe them using a temporary notation: if π is a permutation then $\pi|_{[i \dots j]}$ denotes the subsequence of π whose values comprise the interval $[i \dots j]$. All permutations in \mathcal{C} of length n will belong to one of the following two types:

- permutations belonging to \mathcal{A} ;
- permutations π not belonging to \mathcal{A} which have the property that if k is the minimum value such that $\pi|_{[1 \dots k]} \notin \mathcal{A}$ then $\pi|_{[(k+1) \dots n]} \in \mathcal{B}$.

Consider the collection of permutations not belonging to \mathcal{A} but which have the property that the permutation resulting from the deletion of their maximum symbol does lie in \mathcal{A} . If we define \hat{a}_n to be the number of permutations of this type of length n then it is easy to see that:

$$\hat{a}_n = na_{n-1} - a_n$$

since the first term on the right hand side counts the number of ways of adding a new maximum to a permutation in \mathcal{A} of length $n - 1$ while the second term subtracts the number of ways to do this which still result in a permutation in \mathcal{A} .

The description of the permutations in \mathcal{C} then shows that:

$$c_n = a_n + \sum_{k=0}^n \binom{n}{k} \hat{a}_k b_{n-k}$$

and the theorem follows by comparison of series. ■

So far as we know this is the first appearance of exponential generating functions in pattern class enumeration. Notice from the form of the result that $av(R \oplus S, \mathcal{IW})$ and $av(S \oplus R, \mathcal{IW})$ are equinumerous.

Proposition 3 shows that we can enumerate weak sorting classes using the various techniques that have been developed for ordinary closed classes. We shall begin these enumerative studies by looking at classes with a single basis permutation of length 3 or 4. The length 3 case is virtually trivial. By Lemma 2 we may restrict our attention to the permutations 123, 132, 231, 321 and we have

Proposition 7 The classes $av(123, \mathcal{IW})$, $av(132, \mathcal{IW})$, $av(231, \mathcal{IW})$, $av(321, \mathcal{IW})$ are enumerated by, respectively

1. $a_n = 0$ for all $n \geq 3$,
2. n ,
3. 2^{n-1} ,
4. the Catalan numbers.

For length 4 there is considerably more to do but Theorem 6 handles many of the cases. To within symmetry we have 16 permutations which, for discussion purposes, we have grouped into 4 families:

- (i) 1234;
- (ii) 2134, 1324, 2314, 3124, 3214, 2143;
- (iii) 4231, 3421, 4321;
- (iv) 2341, 2413, 3142, 2431, 3241, 3412.

The single permutation of the first family defines a finite class. The permutations of the second family are all handled by applying Theorem 6 and this gives the following enumerative formulae (valid for all $n \geq 2$):

$$\frac{\begin{array}{cccccc} 2134 & 1324 & 2314 & 3124 & 3214 & 2143 \\ n(n-1) & n(n-1) & n2^{n-2} & n2^{n-2} & \binom{2n-2}{n-1} & n2^{n-1} - 2^n + 2 \end{array}}{}$$

The third family requires that we solve the enumeration problem for the closed classes with bases $\{4231, 4321\}$, $\{3421, 4321\}$, $\{4321\}$. The first of these (sequence A053617 of [11]) has an enumeration scheme in the sense of [14], the second gives the large Schröder numbers [9] and the third has been computed in [7].

The permutations in the last family present a series of different challenges. The easiest are 2341 and 3412. In these cases the classes are (in the notation of the next section) $\mathcal{B}(3, 1)$ and $\mathcal{B}(2, 2)$, and Proposition 20 gives us the recurrence relations $a_n = 3a_{n-1}$ and $a_n = 4a_{n-1} - 2a_{n-2}$ respectively. We treat the others in a series of lemmas.

Lemma 8 *The class $av(2413, \mathcal{IW})$ is enumerated by $\frac{1}{4}(3^n - 2n + 3)$.*

Proof: The upward weak closure of 2413 is the set $\{2413, 4213, 2431, 4231, 4321\}$ but it is convenient instead to enumerate the class whose ordinary basis is $\{3142, 3241, 4132, 4231, 4321\}$ (the inverse class, which is not a weak sorting class). These basis elements tell us that if we have two disjoint descents then the latter lies entirely above the former; they also tell us that we can have at most two immediately adjacent descents.

Now it follows that two disjoint descents must lie in different components and so the indecomposables of the class begin with an increasing sequence, then have at most two down steps and end with an increasing sequence. The number of such having length n is $n2^{n-3}$ if $n \geq 3$. The ordinary generating function of the indecomposables is therefore

$$g(t) = 1 + t + t^2 + \sum_{n=3}^{\infty} n2^{n-3}t^n$$

and the full generating function is $\frac{1}{1-g(t)}$ from which the result follows. ■

Lemma 9 *The class $av(3142, \mathcal{IW})$ is enumerated by $\frac{1}{4}(3^n - 2n + 3)$.*




Figure 1: Indecomposable permutations in $av(2413, \mathcal{IW})$ and $av(3142, \mathcal{IW})$

Proof: Let b_n be the number of indecomposable permutations of length n avoiding the 5 permutations 3142, 3412, 3421, 4312, 4321 of the upward weak closure of 3142. We shall show that $b_n = 2b_{n-1} + 2^{n-3}$ from which follows $b_n = n2^{n-3}$. Then the proof can be completed as in the previous lemma.

First note that, to avoid the permutations 3412, 3421, 4312, 4321, implies that symbol 1 or symbol 2 must occur in the first two positions. Therefore we can divide the indecomposable permutations into subsets (disjoint if $n > 2$) as follows:

1. $F_1 = \{\pi \mid \pi = 1\dots\}$,
2. $F_2 = \{\pi \mid \pi = 2\dots\}$,
3. $S_1 = \{\pi \mid \pi = t1\dots\}$,
4. $S_2 = \{\pi \mid \pi = t2\dots\}$.

If $n > 1$ then, by the indecomposability, F_1 is empty. Furthermore, if the initial symbol 2 is removed from a permutation of F_2 then the result remains indecomposable. Moreover, any indecomposable permutation of the class can be prefaced by a symbol 2 (incrementing the symbols larger than 2) and the result is not only in the class but is indecomposable. This shows that $|F_2| = b_{n-1}$. A similar argument proves that $|S_2| = b_{n-1}$.

Consider now a permutation $t1\dots \in S_1$. Notice that $t \neq 2$ by indecomposability. We shall prove that $t = n$. If not, let s be the rightmost symbol smaller than t and write the permutation as $t1\alpha s\beta$. The avoidance of 3142 shows that α has no symbols larger than t , and β , by definition, has no symbols smaller than t . So β consists precisely of the set $\{t+1, \dots, n\}$ in some order, contradicting indecomposability as $t < n$.

Hence S_1 is the set of permutations $n1\dots$ in the class which is in 1-1 correspondence with permutations of length $n-2$ that avoid 3142, 3412, 3421, 312, 321. These avoidance conditions amount to avoiding 312, 321 alone and so this set has size 2^{n-3} . ■

The equality of the enumerations in the last two lemmas appears to be no more than a coincidence. From the proofs of these lemmas it is not hard to determine the structures of the indecomposable permutations in both cases and we display these in Figure 1.




Figure 2: Structure of a permutation in $\text{av}(2431, \mathcal{IW})$

Lemma 10 For $\text{av}(2431, \mathcal{IW})$ we have the enumeration formula

$$\sum_{k=0}^n \binom{n}{k} f_{n-k}$$

where (f_n) is Fine's sequence A000957 in [11] (see also [6]).

Proof: Let $\mathcal{D} = \text{av}(2431, \mathcal{IW}) = \text{av}(2431, 4231, 4321)$. We shall determine the structure of a permutation $\pi \in \mathcal{D}$. Consider any left to right maximal m of π , that is, any symbol larger than all of its predecessors. Since π avoids 4231 and 4321, the subsequence of those symbols that follow m in π and are also less than m avoids 231 and 321.

Moreover, if $m' < m$ is a right to left maximal preceding m in π then, because π avoids 2431, all the symbols following m and less than m' must occur before any of the symbols following m and greater than m' but less than m .

Let the sequence of left to right maximals in π be m_1, m_2, \dots, m_k , and let B_i for $1 \leq i \leq k$ be the symbols of π to the right of m_i and between m_i and m_{i-1} in value (take $m_0 = 0$ conventionally). Since the m 's are the left to right maximals, they, together with the sets B_i partition the symbols of π . Moreover, the observation above shows that if $i < j$ then all the symbols B_i must precede all of the symbols B_j . Figure 2 illustrates these conditions.

Every permutation of this form belongs to \mathcal{D} and we can construct them all as follows. Choose an increasing sequence m_i from among 1 through n . For each i , let B_i be the set of values strictly between m_{i-1} and m_i and choose a $\{231, 321\}$ -avoiding permutation β_i of B_i . Now merge the sequences $m_1 m_2 \cdots m_k$ and $\beta_1 \beta_2 \cdots \beta_k$ subject only to the condition that m_i precedes β_i for each i . Then the resulting permutation belongs to \mathcal{D} .

We say that m_i is *bound* if B_i is not empty. Otherwise, m_i is *free*. A permutation in \mathcal{D} is *completely bound* if all of its left to right maximals are bound. Consider first the completely bound permutations in \mathcal{D} . We associate to each of these a word in the alphabet a, b, c as follows:

- Each left to right maximal is encoded by the letter c .
- The last symbol of each B_i is encoded by the letter b .
- All remaining symbols are encoded by the letter a .

We note that, read left to right, the number of c 's minus the number of b 's is always non-negative, ends at 0, and that an a may not occur when the count is 0. All sequences meeting these criteria can occur, and

the number of permutations of \mathcal{D} having all left to right maximal bound, corresponding to a sequence containing k a 's is just 2^k (since each block of a 's between two b 's represents, together with the symbol for its final b , a $\{231, 321\}$ -avoiding permutation and there are 2^{j-1} such of length j). So, we can obtain a one to one correspondence between encodings and this subset of \mathcal{D} if we allow the a symbols to be either a_1 or a_2 arbitrarily (or by using a natural encoding of the corresponding B over a two letter alphabet).

This gives a correspondence between the subset of \mathcal{D} in which all left to right maximal are bound, with Motzkin paths where the horizontal steps can have either of 2 types, but may not occur on the axis, and these are enumerated by Fine's sequence [6, 11]. Let f_n denote its n th symbol.

It remains only to insert the free left to right maximal. Now observe that if we take an arbitrary $\pi \in \mathcal{D}$ and delete the free left to right maximal, what remains is indeed a completely bound permutation. Moreover, if we take such a permutation and nominate places in which left to right maximal are to be inserted freely, then there is a unique way to do so. That is, in a permutation belonging to \mathcal{D} of length n we are free to choose the number of free maximal, and their positions, and then the structure of the remaining bound permutation. This gives

$$d_n = \sum_{k=0}^n \binom{n}{k} f_{n-k}$$

as required. ■

Lemma 11 *For $av(3241, \mathcal{IW})$ we have the generating function*

$$\frac{3 - 13t + 2t^2 + 5t\sqrt{1-4t} - \sqrt{1-4t}}{2(1-4t-t^2)}$$

Proof: The WILFPLUS package [13] is able to produce an enumeration scheme for this class from which, in principle, one could obtain the stated generating function. However, we have derived it using techniques developed in [1]. ■

3 Strong sorting classes

For weak sorting classes Proposition 3, Corollary 4 and Corollary 5 describe how the weak basis is related to the ordinary basis. The situation for strong sorting classes is considerably more complex. For example, the direct analogue of Corollary 4 is false since, for example, it would imply $av(321, \mathcal{IS}) = av(321)$; however, $321 \mathcal{I} 3214 \mathcal{S} 3412$ and therefore $3412 \in av(321) \setminus av(321, \mathcal{IS})$. Despite this we shall prove that a strong sorting class with a finite strong basis has a finite ordinary basis and our proof will show how this ordinary basis may be computed from the strong basis.

We begin these investigations by defining three types of operation on permutations τ or their subsequences:

Switch. Exchange two symbols of τ that are currently correctly ordered.

Left. Move a symbol t of τ to the left and insert some new symbol s smaller than t in the original position of t (with appropriate renumbering of all original symbols greater than or equal to s).

Right. Move a symbol t of τ to the right and insert some new symbol larger than t in the original position of t (also with appropriate renumbering of symbols).

It is helpful to represent a permutation τ by its graph (the set of points $(x, \tau(x))$ plotted in the (x, y) -plane) to show the effect of these operations. Our first use of this graphical representation occurs in




Figure 3: The operations switch, left, and right

Figure 3 which shows the effect of a single operation. We shall make heavier use of these diagrams in the proof of Theorem 14.

Suppose that T is some set of permutations. Then T is said to be *complete* if, for any $\tau \in T$, applying any of the types of operation **switch**, **left**, or **right** to τ results in a permutation that contains some permutation in T as a subpermutation.

Proposition 12 *T is complete if and only if $av(T)$ is a strong sorting class.*

Proof: To begin with, assume that T is complete. By definition, $av(T)$ is closed under taking subpermutations and so we must prove that it is also closed downwards in the strong order; in other words, we must prove

$$\sigma \in av(T) \text{ and } \pi \mathcal{S} \sigma \implies \pi \in av(T)$$

and it is clearly sufficient to prove this in the case that π and σ differ by an exchange.

So let $\sigma \in av(T)$ and $\pi \mathcal{S} \sigma$ where π and σ differ by an exchange. For a contradiction suppose that π contains a subsequence $p_1 p_2 \dots$ order isomorphic to an element of T . Now σ and π differ only in that two symbols properly ordered in π are in the other order within σ . If neither of these two swapped symbols are among $p_1 p_2 \dots$ then σ also contains this subsequence, and this is impossible. If both of the swapped symbols are among $p_1 p_2 \dots$ then σ contains a subsequence obtainable from $p_1 p_2 \dots$ by swapping two symbols currently in the right order. But a **switch** operation on an element of T results in a permutation that involves an element of T , so this is also impossible.

If only one of the swapped symbols (p say) is among $p_1 p_2 \dots$ and the other symbol is, say, q then consider the subsequence ξ of σ on the symbols p_1, p_2, \dots, q . If, in π , q was to the left of p then we must have $q < p$. But that means that ξ has been obtained from $p_1 p_2 \dots$ by a **left** operation. Similarly if, in π , q was to the right of p then we must have $q > p$ and ξ has been obtained from $p_1 p_2 \dots$ by a **right** operation. In either case, the completeness property tells us that ξ involves an element of T which is impossible.

For the converse, assume that $av(T)$ is a strong sorting class. Let τ be an arbitrary element of T and suppose that τ^* is the result of applying a **switch**, **left**, or **right** operation to τ . Since strong sorting classes are lower ideals in the order \mathcal{IS} , $\tau^* \notin av(T)$. Hence τ^* involves an element of T and therefore T is complete. ■

Now suppose that X is a strong sorting class with strong basis R . Let $c(R)$ denote the ordinary basis of

X . Our aim is to describe $c(R)$ in terms of R . Let \bar{X} denote the complement of X . Then, by definition

$$\bar{X} = \{\theta \mid \rho \mathcal{IS} \theta \text{ for some } \rho \in R\}.$$

Also, by definition, $c(R)$ is the set of minimal permutations in \bar{X} (minimal with respect to \mathcal{I}). The following result shows that $c(R)$ can be constructed from R by using **switch**, **left**, and **right** operations.

Lemma 13 *Let $\theta \in c(R)$. Then there exists a sequence of permutations*

$$\theta_0, \theta_1, \dots, \theta_k = \theta$$

where $\theta_0 \in R$, each $\theta_i \in c(R)$, and each θ_i is obtained from θ_{i-1} by a **switch**, **left**, or **right** operation. Furthermore, in any sequence beginning at a permutation of R and ending at θ where each term arises from the previous one by a **switch**, **left**, or **right** operation, all permutations in the sequence are in $c(R)$.

Proof: We shall prove the first part of the lemma by induction over \bar{X} with respect to the order \mathcal{IS} . If θ happens to be minimal under \mathcal{IS} then, by definition, $\theta \in R$ and the result is vacuously true. This establishes the base of the induction and we now take θ to be non-minimal under \mathcal{IS} . In that case there exists some $\theta' \in \bar{X}$ with $\theta' \mathcal{IS} \theta$ where this relation between θ' and θ is a covering relation. Since θ is a minimal element for the order \mathcal{I} we cannot have $\theta' \mathcal{I} \theta$ and so we have $\theta' \mathcal{S} \theta$; furthermore, θ can be obtained from θ' by a **switch** operation (exchanging the symbols a and b say).

If $\theta' \in c(R)$ then we can conclude the proof by induction; therefore assume that $\theta' \notin c(R)$. Then there is some permutation $\theta'' \in \bar{X}$ with $\theta'' \mathcal{I}_0 \theta'$; in other words, θ' has been obtained from θ'' by inserting a new symbol c (with appropriate renumbering). If c is neither a nor b then we can interchange the **switch** of a with b , and the insertion of c , to obtain θ by first switching a and b and then inserting c . However, that is impossible since θ is a minimal element of \bar{X} under involvement.

It is now easy to see that, if $c = a$, then θ is formed from θ'' by a **left** operation while, if $c = b$, then θ is formed from θ'' by a **right** operation.

If $\theta'' \in c(R)$ then, again, we can conclude the proof by induction. Hence, for a final contradiction, we shall assume that $\theta'' \notin c(R)$. In that case there is some $\theta''' \in \bar{X}$ with $\theta''' \mathcal{I}_0 \theta''$ and θ'' is the result of inserting some new symbol d into θ''' . If d is neither a nor b then we can obtain θ from θ''' by an appropriate **left** or **right** operation followed by an insertion of d and, as before, this is impossible by the minimality of θ .

Therefore $\{c, d\} = \{a, b\}$ (or, more precisely, the symbols that have been inserted to form θ' from θ''' become a, b after renumbering) and now we can obtain θ from θ''' by inserting a and b directly into their proper places within θ . Again this implies that θ is not minimal and the proof of the first part is complete.

For the second part, suppose we have a sequence of permutations beginning at a permutation of R and ending at θ each being generated from its predecessor by a **switch**, **left**, or **right** operation. Let ϕ be a permutation in this sequence that is not minimal under involvement because it has some subpermutation $\phi' \in \bar{X}$. The **switch**, **left**, and **right** operations that transform ϕ into θ also transform ϕ' and preserve the involvement property. Ultimately, this contradicts the minimality of θ . ■

This lemma indicates how $c(R)$ can be computed from R using a breadth-first search strategy. We begin from R itself and apply **switch**, **left**, and **right** operations discarding any results that contain previously found permutations as subpermutations; and we continue using any new permutations found. We generate new permutations in order of length (by applying the operations to the smallest permutations first, and applying **switch** operations before **left** and **right** operations). Clearly this process will examine and not discard every \mathcal{I} -minimal permutation of \bar{X} . On the other hand, any permutation σ which is not \mathcal{I} -minimal will contain a (shorter) \mathcal{I} -minimal permutation τ which, by Lemma 13 and the choice of search strategy, will have been examined before σ (and not discarded). By virtue of the presence of τ , σ will be discarded. Once no new permutations can be generated we will have found a complete set and, by Lemma 13, this will be $c(R)$. Our next result shows that this process terminates if R is finite.




Figure 4: A switch operation followed by a left operation

Theorem 14 *Let X be a strong sorting class with strong basis R and suppose that R is finite. Then $c(R)$ is also finite.*

Proof: We shall be relying on Lemma 13 which proves that every permutation $\theta \in c(R)$ can be constructed from some permutation in R by a sequence of **switch**, **left**, and **right** operations. In the first part of the proof we shall show that θ can be constructed by a sequence in which all the **left** operations precede all the **right** operations and, in turn, all the **right** operations precede all the **switch** operations. The graphical representations of **switch**, **left**, and **right** introduced previously will be used extensively.

Suppose in the sequence of operations that has realised θ we have a **switch** operation followed by a **left** operation. If the **left** operation was applied to neither of the two symbols that took part in the **switch** operation then it is evident that the same effect can be achieved by a **left** operation followed by a **switch**. However, if the **left** operation was applied to one of the two switched symbols, we must argue more carefully. Diagrammatically we have one of four different cases as shown in Figure 4. Each of these cases can be modified as shown in Figure 5 so that the same effect is obtained by a **left** operation followed by a **switch** operation.

A similar argument shows that any **switch** operation followed by a **right** operation may also be replaced by a **right** operation followed by a **switch**. Therefore the sequence of operations may be assumed to have all the **switch** operations at the end.

Now suppose that in the realisation of θ there is a **right** operation followed by a **left** operation. If the two symbols generated by the **right** operation are not affected by the **left** operation then these two operations can obviously be interchanged. In the contrary case there are, again, four cases as shown in Figure 6. The second and fourth of these are impossible because their result is not a minimal permutation: they each involve a permutation arising from a different **right** operation on the initial configuration. The first and third can be achieved by a **left** operation then a **right** operation; the intermediate configurations are shown in Figure 7.

Now suppose that $\theta \in c(R)$. We take a sequence of **left**, **right** and **switch** operations that generate θ from some $\rho \in R$. By the results above we may assume that all the **left** operations are applied first, followed by all the **right** operations, and finally the **switch** operations. We can regard each **left** operation as one which splits a point of the diagram into two, moves one of them to the left and the other one down. If we




Figure 5: The same effect with left followed by switch




Figure 6: right followed by left: different cases (two impossible)




Figure 7: The same effect with left followed by right




Figure 8: Two left operations

have two **left** operations, the second of which splits one of the points created by the first **left** (as in Figure 8) the result is not minimal since it involves a permutation formed by performing one **left** operation only. On the other hand two **left** operations that are not so linked can be commuted. Thus, no new point gets split by another **left** operation and so the number of **left** operations cannot be more than the original number of points present. Hence the series of **left** operations cannot increase the length by more than a factor of 2. The same is true of the **right** operations and so $|\theta| \leq 4|\rho|$ which completes the proof. ■

We turn now to the enumeration problem for strong sorting classes whose strong basis is finite and give some sample results. Our general method is to first determine the ordinary basis of the class by the process described above and use our experience in closed class enumeration.

As an example of a fairly typical situation we note that

$$c(\{4231\}) = \{4231, 4321, 35142, 45312, 42513, 45132, 35412, \\ 45213, 43512, 456123, 351624, 451623, 356124\}.$$

The next two results summarise the enumerations of all strong sorting classes with a single strong basis permutation of length 3 or 4 (omitting trivial cases or cases that follow from symmetry).

Proposition 15 *For a single strong basis permutation of length 3*

<i>Basis permutation</i>	<i>Ordinary basis</i>	<i>Enumeration</i>
312	{321, 312}	2^{n-1}
321	{321, 3412}	$a_n = 3a_{n-1} - a_{n-2}$ for $n \geq 3$

Proof: The ordinary basis can be confirmed using Proposition 12 and the enumerations are well-known. ■

Proposition 16 *For a single strong basis permutation of length 4*

<i>Name</i>	<i>Basis permutation</i>	<i>Enumeration</i>
<i>I</i>	1234	0 for $n \geq 4$
<i>II</i>	1243	6 for $n \geq 4$
<i>III</i>	1324	4 for $n \geq 4$
<i>IV</i>	1342	$3 \times 2^{n-2}$ for $n \geq 3$
<i>V</i>	1432	$a_n = 3a_{n-1} - a_{n-2}$ for $n \geq 4$
<i>VI</i>	2143	$a_n = 4n - 6$ for $n \geq 2$
<i>VII</i>	2341	$2 \times 3^{n-2}$ for $n \geq 2$
<i>VIII</i>	2413	$a_n = 3a_{n-1} - 2a_{n-2} + 2a_{n-3}$ for $n \geq 4$
<i>IX</i>	2431	$a_n = 4a_{n-1} - 3a_{n-2} + 2a_{n-3}$ for $n \geq 4$
<i>X</i>	3412	$a_n = 4a_{n-1} - 2a_{n-2}$ for $n \geq 3$
<i>XI</i>	3421	$(4^n + 2)/3$ for $n \geq 2$
<i>XII</i>	4231	$a_n = 4a_{n-1} - 2a_{n-2} + 4a_{n-3} - a_{n-5}$ for $n \geq 6$
<i>XIII</i>	4321	$a_n = 4a_{n-1} + a_{n-2} + a_{n-3} - 4$ for $n \geq 5$

Proof: We give a sketch of the proof of the last of these only. The form of the other proofs is similar although the details vary considerably. First of all we use the completion process to determine the basis of $av(4321, \mathcal{LS})$. This turns out to be

$$\{4321, 45132, 45231, 35412, 53412, 45213, 43512, 45312, 456123, 451623, 356124\}.$$

Next we observe that the basis elements 4321, 45132, 45231, 45213, 45312, 456123 show every permutation of the class has a 1 or 2 or 3 in the first 3 places. Denote the number of permutations of length n in the class by a_n . In the following case analysis we use the letter c to stand for any symbol larger than 2, and the letter d for any symbol larger than 3.

The situation for the permutations that have 1 or 2 in their first or second positions is summarised by

Type	Enumeration	Explanation
1α	a_{n-1}	α can be arbitrary
2α	a_{n-1}	α can be arbitrary
$c1\alpha$	$a_{n-1} - a_{n-2}$	$c\alpha$ is arbitrary but cannot start with 2
$c2\alpha$	$a_{n-1} - a_{n-2}$	$c\alpha$ is arbitrary but cannot start with 1

From now on we assume the first two places do not contain a 1 or a 2. The next cases are those where 3 also does not occur in the first two positions but one of 1, 2, 3 is in the third position. Their forms are as follows

Type	Enumeration	Explanation
$dd1\alpha$	$2a_{n-3} - 2$	Discussed below
$dd2\alpha$	0	Uses 4321, 45213 and 45231
$dd3\alpha$	0	Uses 4321 and 45312

With symbol 3 in second place we have the cases:

Type	Enumeration	Explanation
$d31\alpha$	$a_{n-2} - a_{n-3}$	By removing 1, in correspondence with the type $c2\alpha$
$d32\alpha$	0	Uses 4321
$d3d\alpha$	0	Uses 4321, 43512 and 53412

With symbol 3 in first place we have the cases:

Type	Enumeration	Explanation
$3d1\alpha$	$a_{n-2} - a_{n-3}$	By removing 1, in correspondence with the type $c2\alpha$
$3d2\alpha$	$a_{n-2} - a_{n-3}$	By removing 2, in correspondence with the type $c2\alpha$
$3dd\alpha$	$2a_{n-3} - 2$	Discussed below

The explanations above are straightforward except for the two where we promised further discussion. For the first of these (the type $dd1\alpha$) we can prove (by a quite lengthy case by case examination whose

details we omit) that α starts with 2. Let b_n be the number of permutations of this type. By removing the symbol 2 we obtain a correspondence with the type $cc1\alpha$ of length $n - 1$. The latter sequences have one of the forms $3d1\alpha$ ($a_{n-3} - a_{n-4}$ of them), $d31\alpha$ (also $a_{n-3} - a_{n-4}$ of them), or $dd1\alpha$ (b_{n-1} of them). Hence $b_n = b_{n-1} + 2(a_{n-3} - a_{n-4})$. Iterating this recurrence leads to $b_n = b_5 + 2a_{n-3} - 2a_2$ and since $b_5 = a_2 = 2$ the required result follows. The second case where further discussion was promised is the $3dd\alpha$ case. Here we can prove that the sequences are of the form $34d\alpha$ and then we argue in a similar way.

Adding together all these contributions we obtain $a_n = 4a_{n-1} + a_{n-2} + a_{n-3} - 4$. ■

The tenor of the above results hints that the theory of strong sorting classes is going to be more complex than that for weak sorting classes. However, in the remainder of this section we give some compensatory results which go some way to proving that it may actually be *less* complex.

Consider the following family of closed classes. The closed class $\mathcal{B}(r, s)$ is defined by the $r!s!$ (ordinary) basis permutations $\beta\alpha$ where $|\beta| = r, |\alpha| = s$ and every symbol of β is greater than every symbol of α . It follows directly from the definition that, for a permutation π of length n to be a member of $\mathcal{B}(r, s)$, there must not exist subsets $I, J \subseteq \{1, \dots, n\}$ such that $|I| \geq r, |J| \geq s, I < J$ and $\pi(I) > \pi(J)$. Two other readily checked properties are

$$\mathcal{B}(r, s)^* = \mathcal{B}(r, s)^{-1} = \mathcal{B}(s, r).$$

As a first application of Proposition 12 we have

Lemma 17 $\mathcal{B}(r, s)$ is a strong sorting class. Indeed, if

$$\theta_{rs} = s + 1, s + 2, \dots, s + r, 1, 2, \dots, s$$

then

$$\mathcal{B}(r, s) = \text{av}(\theta_{rs}, \mathcal{IS}).$$

Proof: It is readily checked that the basis of $\mathcal{B}(r, s)$ is complete so the first part follows from Proposition 12. For the second part we note that, as $\mathcal{B}(r, s)$ is a strong sorting class not containing θ_{rs} (which is one of its basis permutations), $\mathcal{B}(r, s) \subseteq \text{av}(\theta_{rs}, \mathcal{IS})$. On the other hand every basis permutation β of $\mathcal{B}(r, s)$ satisfies $\theta_{rs} \mathcal{S} \beta$ and hence cannot be involved in any permutation of $\text{av}(\theta_{rs}, \mathcal{IS})$; so $\text{av}(\theta_{rs}, \mathcal{IS}) \subseteq \mathcal{B}(r, s)$. ■

The importance of the strong sorting classes $\mathcal{B}(r, s)$ stems from

Proposition 18 Every proper strong sorting class is contained in some $\mathcal{B}(r, r)$.

Proof: Let X be a strong sorting class contained in no $\mathcal{B}(r, r)$. Then X contains permutations of the form $\beta\alpha$ where $\beta > \alpha$ and $|\alpha| = |\beta| = r$ for all values of r . But, if X contains say $b_1 \dots b_r a_1 \dots a_r$ with all $b_i > a_j$, then by a series of exchanges of the form $b_i \leftrightarrow a_j$ we can produce a permutation with any rearrangement of $a_1 \dots a_r$ in the first r positions. Thus X contains every permutation of length r and so contains every permutation. ■

This proposition indicates that the classes $\mathcal{B}(r, s)$ are going to be fundamental in the understanding of strong sorting classes. Obviously, $\mathcal{B}(1, 1)$ consists only of identity permutations. The first non-trivial cases are $\mathcal{B}(1, 2)$ and $\mathcal{B}(2, 1)$ whose structure is given next. Subsequently, in Theorem 23, we shall give a complete description of the classes $\mathcal{B}(r, s)$.

Lemma 19 $\mathcal{B}(1, 2)$ consists of permutations whose cycle structure is

$$(1, 2, \dots, k_1)(k_1 + 1, k_1 + 2, \dots, k_2)(k_2 + 1 \dots) \dots$$

and $\mathcal{B}(2, 1)$ is the class of their inverses.

Proof: $\mathcal{B}(1, 2)$ is the class of $\{321, 312\}$ -avoiding permutations whose structure is well-known. The second statement follows from $\mathcal{B}(2, 1) = \mathcal{B}(1, 2)^{-1}$. ■

The classes $\mathcal{B}(r, s)$ were also defined (somewhat differently) by Mansour and Vainshtein [10] who enumerated them by generating functions. The following result gives an elementary method of enumerating them.

Proposition 20 *Let x_n be the number of permutations of length n in $\mathcal{B}(r, s)$. Then*

$$x_n = rsx_{n-1} - 2! \binom{r}{2} \binom{s}{2} x_{n-2} + 3! \binom{r}{3} \binom{s}{3} x_{n-3} - \dots$$

Proof: Suppose we have a permutation of $\{1, 2, \dots, n\} \setminus \{t\}$ where t is one of $\{n, n-1, \dots, n-r+1\}$ and that the permutation is in (i.e. order isomorphic to a permutation of) $\mathcal{B}(r, s)$. If we insert the symbol t anywhere within the final s symbols of this permutation we cannot introduce a subpermutation isomorphic to a basis permutation of $\mathcal{B}(r, s)$, so the result is still in $\mathcal{B}(r, s)$.

Now consider the possible forms of a permutation of length n in $\mathcal{B}(r, s)$. Such a permutation must have at least one of the r largest symbols somewhere within its last s positions. The choice of the value of this symbol together with its positions, and the results of the previous paragraph, would appear to give rsx_{n-1} permutations in $\mathcal{B}(r, s)$ of length n . However, this overcounts the permutations which have two or more of their r largest symbols in their final s positions. So we seem to have $rsx_n - 2! \binom{r}{2} \binom{s}{2} x_{n-2}$ permutations. However, this undercounts by $3! \binom{r}{3} \binom{s}{3} x_{n-3}$ the permutations with three or more of their r largest symbols in their final s positions. Continuing by inclusion-exclusion we obtain the formula. ■

In [2] it was observed that, for closed classes \mathcal{X}, \mathcal{Y} , the set of permutation products

$$\mathcal{X} \circ \mathcal{Y} = \{\alpha \circ \beta \mid \alpha \in \mathcal{X}, \beta \in \mathcal{Y}\}$$

was also a closed class. A similar result holds for strong sorting classes.

Proposition 21 *If \mathcal{X} and \mathcal{Y} are strong sorting classes so also is $\mathcal{X} \circ \mathcal{Y}$.*

Proof: A class \mathcal{Z} is closed under the strong order if and only if for all $\zeta \in \mathcal{Z}$ and all i, j with $i < j$ and $i^\zeta > j^\zeta$ we have (using cycle notation for transpositions)

$$(i, j) \circ \zeta \in \mathcal{Z}.$$

Now let $\alpha \in \mathcal{X}$ and $\beta \in \mathcal{Y}$ and put $\gamma = \alpha \circ \beta$. Suppose $i < j$ and $i^\gamma > j^\gamma$. Then $(i, j) \circ \gamma = (i, j) \circ \alpha \circ \beta$. Either $i^\alpha > j^\alpha$ in which case $(i, j) \circ \alpha \in \mathcal{X}$ or $i^\alpha < j^\alpha$ in which case

$$(i, j) \circ \alpha \circ \beta = \alpha \circ \alpha^{-1} \circ (i, j) \circ \alpha \circ \beta = \alpha \circ (i^\alpha, j^\alpha) \circ \beta$$

and $(i^\alpha, j^\alpha) \circ \beta \in \mathcal{Y}$ since $i^{\alpha\beta} > j^{\alpha\beta}$. ■

Lemma 22 $\mathcal{B}(1, q) \supseteq \mathcal{B}(1, 2)^{q-1}$ and $\mathcal{B}(p, 1) \supseteq \mathcal{B}(2, 1)^{p-1}$.

Proof: If the first part were false there would be some basis element of $\mathcal{B}(1, q)$ which was expressible as a product of $q-1$ elements of $\mathcal{B}(1, 2)$. Such a basis element has length $q+1$ and maps 1 to $q+1$ (i.e. as a sequence it begins with $q+1$). However each element of $\mathcal{B}(1, 2)$ maps symbols t either to $t+1$ or to a smaller symbol (Lemma 19) and so a product of $q-1$ of them cannot map 1 to $q+1$. The second part follows by taking inverses. ■

Theorem 23

$$\mathcal{B}(p, q) = \mathcal{B}(2, 1)^{p-1} \circ \mathcal{B}(1, 2)^{q-1}$$

and

$$\mathcal{B}(p, q) \circ \mathcal{B}(r, s) = \mathcal{B}(p+r-1, q+s-1)$$

Proof: We prove a series of results from which we then deduce the theorem.

$$A: (X \circ Y)^* = X^* \circ Y^*.$$

If α, β have length n and ρ is the reverse identity permutation then

$$\begin{aligned} (\alpha \circ \beta)^* &= \rho \circ \alpha \circ \beta \circ \rho \\ &= \rho \circ \alpha \circ \rho \circ \rho \circ \beta \circ \rho \\ &= \alpha^* \circ \beta^*. \end{aligned}$$

$$B: \mathcal{B}(p, q) \supseteq \mathcal{B}(p, 1) \circ \mathcal{B}(1, q).$$

If this is false there would exist a permutation $\pi = \alpha\beta$ with $\alpha \in \mathcal{B}(p, 1), \beta \in \mathcal{B}(1, q)$ and sets I, J with $|I| \geq p, |J| \geq q, I < J$ and $\pi(I) > \pi(J)$. Then, as $\beta(\alpha(I)) > \beta(\alpha(J))$ we have

$$\beta(r) > \beta(s)$$

for all $r \in \alpha(I), s \in \alpha(J)$. For each fixed $r \in \alpha(I), r < \alpha(J)$ is impossible since $\beta \in \mathcal{B}(1, q)$. Thus, for all $r \in \alpha(I)$, we have $r > s_0$ where s_0 is the minimal value in $\alpha(J)$. But now, writing $j = \alpha^{-1}(s_0)$ we have $I < j$ and $\alpha(I) > \alpha(j)$ contradicting that $\alpha \in \mathcal{B}(p, 1)$.

$$C: \mathcal{B}(p, q) \subseteq \mathcal{B}(2, 1) \circ \mathcal{B}(p-1, q).$$

We say that a subsequence of length $r + s$ is of type (r, s) if its initial r symbols are all greater than the final s symbols. Let $\sigma \in \mathcal{B}(p, q)$. By definition σ has no subsequences of type (p, q) . A subsequence γ of σ is defined to be *critical* if

1. $|\gamma| = p - 1$, and
2. the set $S(\gamma)$ of symbols that follow γ and are less than every symbol of γ has size at least q .

The critical subsequences γ together with the q -symbol subsequences of $S(\gamma)$ comprise all the subsequences of σ of type $(p-1, q)$.

Notice that two distinct critical sequences γ_1, γ_2 cannot have the same last symbol. For then $\gamma_1 \cup \gamma_2$ would have size at least p and the minimal symbol of this subsequence would be one of $\min(\gamma_1)$ and $\min(\gamma_2)$ and so this subsequence, together with either $S(\gamma_1)$ or $S(\gamma_2)$ would contain a subsequence of type (p, q) . We can therefore order the critical sequences by the position in σ of their final symbols.

Let γ be any critical sequence with final symbol g and let $x_r, x_{r-1}, \dots, x_q, \dots, x_1$ be the symbols of $S(\gamma)$ in order of their occurrence in σ . There can be no symbols of σ between g and x_q except for x_r, \dots, x_{q+1} . For if there were such a symbol it could not be smaller than $\min(\gamma)$ (or, by definition, it would be in $S(\gamma)$) and, if it were larger than $\min(\gamma)$ then together with γ and x_q, \dots, x_1 we would have a subsequence of type (p, q) . Notice now that if the segment $gx_r x_{r-1} \dots x_q$ was replaced by $x_r x_{r-1} \dots x_q g$ then we would destroy all the subsequences of type $(p-1, q)$ that stemmed from γ ; moreover no further subsequences of type $(p-1, q)$ would be introduced. This replacement can be effected by pre-multiplication of σ by a cycle $(i, i+1, \dots, j)$ where i and j are the initial and final positions of the segment $gx_r x_{r-1} \dots x_q$.

Now, the next critical sequence γ' (which ends after g) cannot have final symbol one of x_r, x_{r-1}, \dots, x_q . If it did then $\gamma \cup \gamma'$ together with $S(\gamma')$ would contain a subsequence of type (p, q) . Therefore the next critical sequence has a final symbol which occurs after x_q . That shows that the subsequences of type $(p-1, q)$ that stem from γ' can be destroyed by pre-multiplication by a similar cycle disjoint from the one defined above.

Hence all the subsequences of type $(p-1, q)$ can be destroyed by pre-multiplication by a product of disjoint cycles of the form given above. But such a product is, by Lemma 19, a permutation $\alpha \in \mathcal{B}(1, 2)$. So there exists some $\alpha \in \mathcal{B}(1, 2)$ for which $\alpha\sigma \in \mathcal{B}(p-1, q)$ and, as $\mathcal{B}(1, 2)^{-1} = \mathcal{B}(1, 2)$ we have $\sigma \in \mathcal{B}(2, 1) \circ \mathcal{B}(p-1, q)$ as required.

D: $\mathcal{B}(p, q) = \mathcal{B}(p, 1) \circ \mathcal{B}(1, q) = \mathcal{B}(2, 1)^{p-1} \circ \mathcal{B}(1, 2)^{q-1}$.

By repeated application of C we have

$$\mathcal{B}(p, q) \subseteq \mathcal{B}(2, 1)^{p-1} \circ \mathcal{B}(1, q) \subseteq \mathcal{B}(p, 1) \circ \mathcal{B}(1, q)$$

(the second inclusion following from Lemma 22) and the result follows from B.

E: $\mathcal{B}(p, 1) \circ \mathcal{B}(r, 1) = \mathcal{B}(p + r - 1, 1)$ and $\mathcal{B}(1, q) \circ \mathcal{B}(1, s) = \mathcal{B}(1, q + s - 1)$.

This follows immediately from D since $\mathcal{B}(1, 1)$ is the class of identity permutations.

F: $\mathcal{B}(p, 1) \circ \mathcal{B}(1, q) = \mathcal{B}(1, q) \circ \mathcal{B}(p, 1)$.

The left hand side is $\mathcal{B}(p, q)$ (by D) and the right hand side is

$$\begin{aligned} \mathcal{B}(1, q) \circ \mathcal{B}(p, 1) &= \mathcal{B}(q, 1)^* \circ \mathcal{B}(1, p)^* \\ &= (\mathcal{B}(q, 1) \circ \mathcal{B}(1, p))^* \\ &= \mathcal{B}(q, p)^* \\ &= \mathcal{B}(p, q). \end{aligned}$$

We can now complete the proof since

$$\begin{aligned} \mathcal{B}(p, q) \circ \mathcal{B}(r, s) &= \mathcal{B}(p, 1) \circ \mathcal{B}(1, q) \circ \mathcal{B}(r, 1) \circ \mathcal{B}(1, s) \\ &= \mathcal{B}(p, 1) \circ \mathcal{B}(r, 1) \circ \mathcal{B}(1, q) \circ \mathcal{B}(1, s) \\ &= \mathcal{B}(p + r - 1, 1) \circ \mathcal{B}(1, q + s - 1) \\ &= \mathcal{B}(p + r - 1, q + s - 1). \end{aligned}$$

■

Finally we mention another respect in which strong sorting classes are more tractable than weak sorting classes. It is no accident that the enumerations in Propositions 15 and 16 were all rational since in [1] techniques are given to show that every finitely based subclass of $\mathcal{B}(r, s)$ has a rational generating function. Consequently, using Theorem 14 we have

Theorem 24 *Every strong sorting class whose strong basis is finite has a rational generating function.*

Acknowledgement We made extensive use of the GAP system [5] in deriving some of the enumerative results. We also thank the referee whose very careful reading of the paper resulted in many improvements.

References

- [1] M. H. Albert, S. A. Linton, N. Ruškuc: The insertion encoding of permutations, Department of Computer Science, University of Otago, Technical Report OUCS-2005-05 (see <http://www.cs.otago.ac.nz/research/publications/oucs-2005-05.pdf>).
- [2] M. H. Albert, R. E. L. Aldred, M. D. Atkinson, H. van Ditmarsch, C. C. Handley, D. A. Holton, D. J. McCaughan: Compositions of pattern restricted sets of permutations, in preparation.
- [3] M. D. Atkinson, D. Tulley, M. J. Livesey: Permutations generated by token passing in graphs, Theoretical Computer Science 178 (1997), 103–118.
- [4] M. D. Atkinson, T. Stitt: Restricted permutations and the wreath product, Discrete Math. 259 (2002), 19–36.

- [5] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4*; 2004, (<http://www.gap-system.org>).
- [6] E. Deutsch, L. Shapiro: A survey of the Fine numbers, *Discrete Math.* 241 (2001), 1–3.
- [7] I. M. Gessel: Symmetric functions and P -recursiveness, *J. Combin. Theory Ser. A* 53 (1990), 257–285.
- [8] D. E. Knuth: *Fundamental Algorithms, The Art of Computer Programming* Vol. 1 (Second Edition), Addison-Wesley, Reading, Mass. (1973).
- [9] D. Kremer: Permutations with forbidden subsequences and a generalised Schröder number, *Discrete Math.* 218 (2000), 121–130.
- [10] T. Mansour, A. Vainshtein: Avoiding maximal parabolic subgroups of S_k , *Discrete Mathematics and Theoretical Computer Science* 4 (2000), 67–77.
- [11] N. J. A. Sloane: Superseeker, superseeker@research.att.com.
- [12] *Special issue on permutation patterns*, *Electron. J. Combin.* 9(2) (2003), editors M. D. Atkinson, D. A. Holton.
- [13] V. Vatter: WILFPLUS, Maple package available at <http://www.math.rutgers.edu/~vatter/programs/wilfplus/index.html>.
- [14] D. Zeilberger: Enumeration Schemes, and More Importantly, Their Automatic Generation, *Annals of Combinatorics* 2 (1998), 185–195.